

A Comment on “Weak Phase γ using Isospin Analysis and Time-dependent Asymmetry in $B^0 \rightarrow K_S \pi^+ \pi^-$ ”¹Michael Gronau²*Enrico Fermi Institute and Department of Physics
University of Chicago, Chicago, Illinois 60637*

We comment on a recent suggestion for measuring γ in $B^0(t) \rightarrow K_S \pi^+ \pi^-$. A difficulty is pointed out in relating electroweak penguin and tree amplitudes in $B^+ \rightarrow K^0 \pi^+ \pi^0$, which is crucial for an implementation of this method. We clarify the necessary condition for such a relation.

PACS codes: 12.15.Hh, 12.15.Ji, 13.25.Hw, 14.40.Nd

In a recent interesting Letter [1] Nilendra Deshpande, Nita Sinha and Rahul Sinha propose to use $B \rightarrow K \pi \pi$ decays in order to determine the weak phase γ . Their method is based on the CP asymmetry in $B^0(t) \rightarrow K_S \pi^+ \pi^-$, and on an isospin triangle relation among the three amplitudes for $B^+ \rightarrow K^0(\pi^+ \pi^0)_e$, $B^0 \rightarrow K^0(\pi^+ \pi^-)_e$ and $B^0 \rightarrow K^0(\pi^0 \pi^0)_e$, in which the two pions are in an even angular momentum state. A crucial assumption of this method is that electroweak penguin and tree amplitudes contributing to $B^+ \rightarrow K^0(\pi^+ \pi^0)_e$ involve a common strong phase. Such a property was shown to hold in the SU(3) symmetry limit for the $I = 3/2$ amplitude in $B \rightarrow K \pi$ [2, 3], and in the isospin symmetry limit for the $I = 2$ $B \rightarrow \pi \pi$ amplitude [3, 4].

The purpose of this short comment is to clarify the general condition under which tree and electroweak amplitudes in charmless B decays can be related to each other. We will show that this condition is not fulfilled in the case studied in [1].

The effective Hamiltonian describing charmless $\Delta S = 1$ (or $\Delta S = 0$) decays [5] consists of current-current operators Q_1 and Q_2 , QCD penguin operators Q_i , $i = 3 - 6$, and electroweak penguin (EWP) operators Q_i , $i = 7 - 10$. The operators Q_1 and Q_2 , multiplying Wilson coefficients c_1 and c_2 , respectively, and CKM coefficients $V_{ub}^* V_{us}$ (or $V_{ub}^* V_{ud}$), will be named tree operators. EWP operators involve CKM factors $V_{tb}^* V_{ts}$ (or $V_{tb}^* V_{td}$). The EWP operators Q_9 and Q_{10} with the dominant Wilson coefficients, c_9 and c_{10} , have the same (V-A)(V-A) structure as the tree operators, and would have approximately the same matrix elements if they had also identical flavor SU(3) and isospin structure.

In order to find out when such a relation holds, one decomposes all four quark operators of the form $(\bar{b}q_1)(\bar{q}_2q_3)$ transforming as $\mathbf{\bar{3}} \otimes \mathbf{3} \otimes \mathbf{\bar{3}}$ in terms of a sum of $\mathbf{\bar{15}}$, $\mathbf{6}$ and $\mathbf{\bar{3}}$ [6]. The representation $\mathbf{\bar{3}}$ appears both symmetric ($\mathbf{\bar{3}}^{(s)}$), and antisymmetric

¹To be published as a Comment in Physical Review Letters

²Permanent Address: Physics Department, Technion – Israel Institute of Technology, 32000 Haifa, Israel.

$(\overline{\mathbf{3}}^{(a)})$ under the interchange of q_1 and q_3 . The tree and electroweak parts of the $\Delta S = 1$ Hamiltonian are [3]:

$$\mathcal{H}_T = -\frac{G_F}{\sqrt{2}} V_{ub}^* V_{us} \left[\frac{c_1 - c_2}{2} (\overline{\mathbf{3}}_0^{(a)} + \mathbf{6}_1) + \frac{c_1 + c_2}{2} (\overline{\mathbf{15}}_1 + \frac{1}{\sqrt{2}} \overline{\mathbf{15}}_0 - \frac{1}{\sqrt{2}} \overline{\mathbf{3}}_0^{(s)}) \right] , \quad (1)$$

$$\mathcal{H}_{\text{EWP}} = -\frac{G_F}{\sqrt{2}} \frac{3V_{tb}^* V_{ts}}{2} \left[\frac{c_9 - c_{10}}{2} (\frac{1}{3} \overline{\mathbf{3}}_0^{(a)} + \mathbf{6}_1) + \frac{c_9 + c_{10}}{2} (-\overline{\mathbf{15}}_1 - \frac{1}{\sqrt{2}} \overline{\mathbf{15}}_0 - \frac{1}{3\sqrt{2}} \overline{\mathbf{3}}_0^{(s)}) \right] , \quad (2)$$

where subscripts denote the isospin of corresponding operators. Note that both the $\mathbf{6}$ and $\overline{\mathbf{15}}$ operators include a $\Delta I = 1$ component.

Eqs. (1) and (2) imply proportionality relations between EWP and tree operators transforming as $\overline{\mathbf{15}}$ and $\mathbf{6}$ [7]:

$$\mathcal{H}_{\text{EWP}}(\overline{\mathbf{15}}) = -\frac{3}{2} \frac{c_9 + c_{10}}{c_1 + c_2} \frac{V_{tb}^* V_{ts}}{V_{ub}^* V_{us}} \mathcal{H}_T(\overline{\mathbf{15}}) , \quad (3)$$

$$\mathcal{H}_{\text{EWP}}(\mathbf{6}) = \frac{3}{2} \frac{c_9 - c_{10}}{c_1 - c_2} \frac{V_{tb}^* V_{ts}}{V_{ub}^* V_{us}} \mathcal{H}_T(\mathbf{6}) . \quad (4)$$

The two proportionality constants are approximately renormalization scale independent, and are approximately equal in magnitudes but differ in sign [5],

$$\frac{c_9 + c_{10}}{c_1 + c_2} \approx \frac{c_9 - c_{10}}{c_1 - c_2} . \quad (5)$$

Therefore, EWP and tree amplitudes in B decay processes which obtain contributions from either the $\overline{\mathbf{15}}$ or the $\mathbf{6}$ operator, but not from both, are proportional to each other and involve a common strong phase. This property does not hold when the two operators contribute because of the opposite signs in Eqs. (3) and (4). In this case EWP and tree amplitudes involve the sum and difference of two complex amplitudes and have different strong phases.

In the case of $B \rightarrow (K\pi)_{I=3/2}$ [2, 3], the K and π are in an S-wave state, which is symmetric under an interchange of the two SU(3) octets. This state is a pure $\mathbf{27}$. The only SU(3) operator which contributes to this transition is the $\overline{\mathbf{15}}$. Consequently, the EWP and tree amplitudes are proportional to each other in the SU(3) approximation. The same holds true in the isospin symmetry limit for the EWP and tree amplitudes of $B \rightarrow (\pi\pi)_{I=2}$, since only the $\overline{\mathbf{15}}$ contains a $\Delta I = 3/2$ component [3, 4]. On the other hand, in $B^+ \rightarrow K^0(\pi^+\pi^0)_e$ studied in [1] the final state has $I = 3/2$, $S = 1$ and can be in a $\mathbf{27}$ and in a $\mathbf{10}$, to which the $\Delta I = 1$ components of both the $\overline{\mathbf{15}}$ and the $\mathbf{6}$ operators contribute. Hence, the condition for proportional EWP and tree amplitudes and for a common strong phase does not hold. Although this proportionality does not follow from symmetry considerations alone, it would be interesting to study possible dynamical assumptions which can lead to such a situation.

I thank Jonathan Rosner for useful discussions, and I am grateful to the Enrico Fermi Institute at the University of Chicago for its kind hospitality. This work was supported in part by the United States Department of Energy through Grant No. DE FG02 90ER40560.

References

- [1] N. G. Deshpande, N. Sinha and R. Sinha, Phys. Rev. Lett. **90**, 061802 (2003).
- [2] M. Neubert and J. L. Rosner, Phys. Lett. B **441**, 403 (1998).
- [3] M. Gronau, D. Pirjol and T. M. Yan, Phys. Rev. D **60**, 034021 (1999).
- [4] A. J. Buras and R. Fleischer, Eur. Phys. J. C **11**, 93 (1999).
- [5] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [6] D. Zeppenfeld, Zeit. Phys. C **8**, 77 (1981); M. Savage and M. Wise, Phys. Rev. D **39**, 3346 (1989); M. Gronau, O. Hernández, D. London, and J. L. Rosner, Phys. Rev. D **50**, 4529 (1994); Phys. Rev. D **52**, 6374 (1995).
- [7] M. Gronau, Phys. Rev. D **62**, 014031 (2000).